Cialis, Levitra, and Viagra Prior Authorization Request Form



3. Is the PDE-5 inhibitor being prescribed for a

diagnosis of Raynaud's phenomenon?

4. What is the dosing regimen?

5609

Please complete Question 4

□ Yes

Coverage approved
Please complete Question 4

Page 1 of 3

To be completed and signed by the prescriber. To be used only for prescriptions which are to be filled through the Department of Defense (DoD) TRICARE pharmacy program (TPHARM). Express Scripts is the TPHARM contractor for DoD. • The provider may call: 1-866-684-4488 or the completed form may be faxed to: 1-866-684-4477 · The patient may attach the completed form to the prescription and mail it to: Express Scripts, P.O. Box 52150, Phoenix, AZ 85072-9954 or email the form only to: TpharmPA@express-scripts.com Prior authorization criteria and a copy of this form are available at https://rxnet.army.mil/pec/forms_criteria.php.. This prior authorization has no expiration date. Please complete patient and physician information (Please Print) Step Patient Name: Physician Name: Address: Address: Sponsor ID# Phone #: Date of Birth: Secure Fax #: Please indicate which medication is being prescribed:

Cialis Levitra Viagra Step Please consider the following: Patients taking nitrates, either regularly or intermittently, should not receive PDE-5 inhibitors. Patients should be informed of the consequences should they initiate nitrate therapy while taking a PDE-5 inhibitor. Please see product labeling precautions for concurrent use with alpha blockers. Please indicate the patient's gender. Step Please go to Section 1 for Female patients □ Female Please go to Section 2 for Male patients on Page 2 □ Male Section 1 – Female patients □ No 1. Is the PDE-5 inhibitor being prescribed for the ☐ Yes **Proceed to Question 2** treatment of sexual dysfunction? Coverage not approved 2. Is the PDE-5 inhibitor being prescribed for a □ No □ Yes Proceed to Question 3 diagnosis of pulmonary arterial hypertension (PAH)? Coverage approved

Please go to Step 4 on Page 3.

□ No

Coverage not approved

Cialis, Levitra, and Viagra Prior Authorization Request Form



5610

Page 2 of 3

Section 2 - Male patients

1. Is the requested medication Levitra?	☐ Yes Proceed to Question 2	☐ No STOP. Go to Section 3 on Page 3
2. Is the patient 18 years of age or older?	☐ Yes Proceed to Question 3	☐ No Proceed to Question 6
3. Is the patient 40 years of age or older?	☐ Yes Do not submit form. Prior authorization is not required for males 40 years or older.	□ No Proceed to Question 4
4. Is the PDE-5 inhibitor being prescribed for the treatment of erectile dysfunction of organic origin? Organic impotence is considered a consequence of chronic medical conditions that result in impaired arterial blood flow or nerve damage, mixed organic/psychogenic causes, and necessary use of causative medications that cannot be reduced or discontinued. TRICARE regulations specifically exclude coverage of therapies for erectile dysfunction that is not of organic origin.	☐ Yes Coverage approved	□ No Proceed to Question 5
5. Is the indication for the preservation or restoration of erectile function following prostatectomy?	☐ Yes Coverage approved To determine quantity requirements, please complete Question 8	☐ No Proceed to Question 6
6. Is the PDE-5 inhibitor being prescribed for a diagnosis of pulmonary arterial hypertension (PAH)?	☐ Yes Coverage approved To determine quantity requirements, please complete Question 8	☐ No Proceed to Question 7
7. Is the PDE-5 inhibitor being prescribed for a diagnosis of Raynaud's phenomenon?	☐ Yes Coverage approved To determine quantity requirements, please complete Question 8	☐ No Coverage not approved
8. What is the dosing regimen?		

Note: For treatment of erectile dysfunction of organic origin unrelated to therapy for preservation or restoration of erectile function following prostatectomy, coverage is limited to a collective quantity (sildenafil, vardenafil and/or tadalafil combined) of 6 tablets per 30 days or 18 tablets per 90 days.

Please go to Step 4 on Page 3.

Cialis, Levitra, and Viagra Prior Authorization Request Form



Step

5611

Page 3 of 3

Has the patient received a trial of Levitra (vardenafil) and had an inadequate response?	☐ Yes	☐ No Proceed to Question 2
2. Has the patient received a trial of Levitra (vardenafil), but was unable to tolerate it due to adverse effects?	Please sign and date	□ No
	Please sign and date	Proceed to Question 3
3. Is treatment with Levitra (vardenafil) contraindicated for this patient (e.g., due to hypersensitivity)?	□ Yes	□ No
	Please sign and date	Proceed to Question 4
4. Is the requested medication being used for treatment of pulmonary arterial hypertension (PAH)?	□ Yes	□ No
	Please sign and date	Proceed to Question 5
	Please complete Question 7	
5. Is the requested medication being used for preservation/restoration of erectile function after prostatectomy?	□ Yes	□ No
	Please sign and date	Proceed to Question 6
	Please complete Question 7	
6. Is the requested medication being used for treatment of Raynaud's phenomenon?	□ Yes	□ No
	Please sign and date	Coverage not approved
	Please complete Question 7	
7. What is the dosing regimen?		

I certify the above is correct and accurate to the best of my knowledge. Please sign and date:

Prescriber signature

December 2009

Date